

**SOLUTIONS FOR EXTRA ADMISSIONS TEST IN  
MATHEMATICS, COMPUTER SCIENCE AND JOINT SCHOOLS  
DECEMBER 2022**

**A** Note that the probability that the coin does not land on heads is  $1 - \cos^2 \alpha = \sin^2 \alpha$ . The probability of exactly two heads is  $\binom{3}{2} \cos^4 \alpha \sin^2 \alpha$  and the probability of exactly three heads is  $\binom{3}{1} \cos^6 \alpha$ . Simplifying gives

$$3 \cos^4 \alpha \sin^2 \alpha + \cos^6 \alpha = 3(1 - \sin^2 \alpha)^2 \sin^2 \alpha + (1 - \sin^2 \alpha)^3 = 1 - 3 \sin^4 \alpha + 2 \sin^6 \alpha$$

**The answer is (d)**

**B** If we write  $u = e^{-x/2}$  then the given expression is  $u - u^2$ . This quadratic has roots at 0 and 1, and is positive in between. Now as  $x$  varies from 0 to  $\infty$ ,  $u$  will decay exponentially from 1 to 0. Along the way,  $u - u^2$  will start at 0, then vary through positive numbers, first increasing for  $u > 1/2$  then decreasing for  $u < 1/2$  and eventually tending towards 0.


**The answer is (a)**

**C** The left-hand side factorises to give

$$(x - 1)(x - 2) < \frac{(x - 1)}{x}$$

When are the two sides equal? That happens when  $x = 1$  or when  $x(x - 2) = 1$ , that is when  $x = 1 \pm \sqrt{2}$ . Note that one of these roots is negative and the other is positive.


That helps us to draw a graph (it's perhaps useful to note that the right-hand side is  $1 - 1/x$ ).



From the graph, we see that the quadratic is below the reciprocal graph for  $1 - \sqrt{2} < x < 0$  and for  $1 < x < 1 + \sqrt{2}$ .

**The answer is (b)**

**D** The first inequality describes a region between two concentric circles. The second inequality is true if either (case 1)  $x \geq 0$  and  $-x \leq \sqrt{3}y \leq x$ , or (case 2)  $x \leq 0$  and  $x \leq \sqrt{3}y \leq -x$ .



The intersection points between the inner circle and those lines are  $(\sqrt{3}/2, 1/2)$  and reflections, so the acute angle subtended at the origin by the two lines is  $60^\circ$ . We would therefore like one-third of the area between the two circles, which is  $\frac{1}{3}\pi(2^2 - 1^2) = \pi$ .

**The answer is (b)**


**E** The centre of the circle is  $\left(\frac{p}{2}, \frac{q+1}{2}\right)$  and the radius is  $\sqrt{\frac{p^2}{4} + \frac{(q-1)^2}{4}}$ , so the equation of the circle is

$$\left(x - \frac{p}{2}\right)^2 + \left(y - \frac{q+1}{2}\right)^2 = \frac{p^2}{4} + \frac{(q-1)^2}{4}$$

On the line  $y = 0$ , this a quadratic for  $x$ . We can set the discriminant equal to zero for a repeated solution. The quadratic is  $x^2 - px + q = 0$  and the discriminant is  $p^2 - 4q$ .

**The answer is (c)**

**F** We need  $-1 < 1 + x - x^2 < 1$ . Let's find the points where  $1 + x - x^2 = 1$  (that's 0 and 1) and find the points where  $1 + x - x^2 = -1$  (that's -1 and 2). Consider the graph.



We need either  $-1 < x < 0$  or  $1 < x < 2$ .

**The answer is (c)**

**G** All of the options are of the form  $y = kf(x)$  with  $k$  constant so let's try that in the equation. We get  $k\frac{df(x)}{dx}$  on the left and  $2(kf(x))^{1/4}$  on the right. Now  $\frac{df(x)}{dx} = (f(x))^{1/4}$  so we just need  $k = 2k^{1/4}$ . We can rearrange this to get  $k = 2^{4/3}$ .

**The answer is (c)**

**H** Clearly  $f$  is always a positive whole integer from the recursion relations.

First find  $n$  with  $f(n) = 1$ . Because  $f(2n) = f(n)$  and  $f(1) = 1$ , it must be the case that for all powers of 2,  $f(n) = 1$ . On the other hand,  $f(2n+1) = f(n) + f(n+1)$  is definitely at least two, so  $f(n)$  is only equal to 1 on the powers of 2.

Next consider values of  $n$  for which  $f(n) = 2$ . The first equation shows us that if we have  $n$  even and  $f(n) = 2$  then  $f(n/2) = 2$ . Let's suppose instead that  $n$  is odd and write  $n = 2m+1$ . Then we would have  $f(n) = f(m) + f(m+1)$ . The only way for this to be 2 is if the right-hand side is  $1 + 1$ , which only happens if both the consecutive numbers  $m, m+1$  are powers of 2, which only happens for 1, 2. Check that  $f(3) = f(1) + f(2) = 2$ . So the points where  $f(n) = 2$  are precisely those where  $n$  is three times a power of 2 (they're of the form  $3 \times 2^k$ ).

Now finally consider values of  $n$  for which  $f(n) = 3$ . The first equation again tells us that we can multiply any such  $n$  by 2. Let's look for odd solutions. They must come from  $1 + 2$  on the right-hand side of the second equation, which only happens if a power of 2 ( $2^l$  say) is one more or one less than a number of the form  $3 \times 2^k$ . That's a bit unusual though, because both  $2^l$

and  $3 \times 2^k$  are even, unless one of the powers of 2 is equal to 1. So those consecutive numbers must be either 2 and 3 or they could be 3 and 4. This gives  $f(5) = 3$  and  $f(7) = 3$ , but that's it for odd numbers. The solutions to  $f(n) = 3$  are therefore  $5 \times 2^k$  or  $7 \times 2^k$ . None of these are multiples of 35.

**The answer is (a)**

**I** Take two to the power of each side for  $x = (x + a)2^b$

The right-hand side is a line with gradient  $2^b$  and  $y$ -intercept  $a2^b$ . We're looking for positive solutions, and there are two ways this could happen; either  $a2^b$  is larger than zero, but the line grows slower than  $x$ , or  $a2^b$  is negative but the line grows faster than  $x$ . This corresponds to (in the first case)  $a > 0$  and  $b < 0$ , or (in the second case)  $a < 0$  and  $b > 0$ . Either way,  $ab < 0$ . There's only one intersection between two straight lines, if any.

**The answer is (c)**

**J** Consider adding together these integrals to get

$$\int_1^2 \frac{x^2 + x^{-2}}{1 + x^4} dx.$$

Now note that the numerator is  $x^{-2}$  multiplied by the denominator, so this simplifies to

$$\int_1^2 x^{-2} dx = [-x^{-1}]_1^2 = \frac{1}{2}$$

The integral we're asked for is therefore  $\frac{1}{2} - A$ .

**The answer is (e)**