

# EXTRA MATHEMATICS ADMISSIONS TEST

November 2023

Time allowed: 1 hour

Full Name	
UCAS ID	
MAT ID	

This paper contains 10 multiple choice questions.

**Calculators are not permitted.**

For each question on pages 2–11 you will be given **five** possible answers, just one of which is correct. Indicate for each question **A–J** which answer (a), (b), (c), (d), or (e) you think is correct with a tick (✓) in the corresponding column in the table below.

	(a)	(b)	(c)	(d)	(e)
A					
B					
C					
D					
E					
F					
G					
H					
I					
J					

**A.** The function  $p(x) = x^3$  is an example of a polynomial with the property that there is a point on the graph  $y = p(x)$  with zero derivative which is neither a local maximum nor a local minimum. Which one of the following polynomials has the same property?

(a)  $y = x^3 - 3x^2 + x$

(b)  $y = x^3 - 3x^2 + 2x$

(c)  $y = x^3 - 3x^2 + 3x$

(d)  $y = x^3 - 3x^2 + 4x$

(e)  $y = x^3 - 3x^2 + 5x$

Please turn over

**B.** Which of the following numbers is the smallest?

(a)  $\log_{10} \left( \sqrt[10]{10^{11}} \right)$

(b)  $\frac{\pi}{2}$

(c)  $\frac{11}{9}$

(d)  $\sqrt{\frac{3}{2}}$

(e)  $\sqrt{3} \cos(44^\circ)$

Please turn over

C. The sum  $\sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4}$ . It follows that

$$1^3 + 3^3 + 5^3 + 7^3 + \dots + 19^3$$

is equal to

- (a) 19,800
- (b) 19,900
- (c) 20,000
- (d) 20,100
- (e) 20,200

Please turn over

**D.** All even square numbers are multiples of 4. All odd square numbers are one more than a multiple of 4. It follows that the number of positive integer solutions  $(x, y)$  to the equation

$$x^2 + 3y^2 = 4442$$

is

- (a) 0
- (b) 1
- (c) 2
- (d) 3
- (e) 4442

Please turn over

**E.** Let  $d(n)$  be the number of digits in a positive integer  $n$  (with  $n$  written in the usual decimal notation). For example  $d(2) = 1$ ,  $d(103) = 3$ , and  $d(10^6) = 7$ .

Define the sequence  $s_n = (20)^{-d(n)}$ . What is the sum  $\sum_{n=1}^{\infty} s_n$  equal to?

(a)  $\frac{1}{2}$

(b)  $\frac{4}{5}$

(c)  $\frac{9}{10}$

(d) 1

(e)  $\frac{9}{5}$

Please turn over

**F.** For two vectors  $\begin{pmatrix} a \\ b \end{pmatrix}$  and  $\begin{pmatrix} c \\ d \end{pmatrix}$  with integer components (positive or negative or zero), we define the function

$$f\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} c \\ d \end{pmatrix}\right) = \begin{pmatrix} ac + bd \\ ad + bc + 2bd \end{pmatrix}.$$

How many vectors  $\begin{pmatrix} a \\ b \end{pmatrix}$  with integer components are there such that

$$f\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} a \\ b \end{pmatrix}\right) = \begin{pmatrix} 2 \\ 0 \end{pmatrix} \quad ?$$

- (a) 0
- (b) 1
- (c) 2
- (d) 3
- (e) Infinitely many

Please turn over

**G.** For a pair of integers  $x$  and  $y$  with  $x \geq 0$  and  $y > 0$ , we define

$$f(x, y) = \frac{1}{2}(x + y)(x + y + 1) + y$$

What is the set of possible values that  $f(x, y)$  can take?

- (a) All positive integers.
- (b) All positive even integers.
- (c) All positive integers except for odd prime numbers.
- (d) All positive integers that are triangular numbers (those which are the sum of the first  $k$  positive integers for some  $k \geq 1$ ).
- (e) All positive integers except for the triangular numbers.

Please turn over



**H.** Let  $p(x) = 2x^4 - 3x^3 - 5x^2 + 2x + 2$ . Given that the  $y = mx$ , with  $m$  a real number, crosses the curve  $y = p(x)$  at four distinct points, let the  $x$ -coordinates of those points be  $x_1, x_2, x_3$ , and  $x_4$ . The product  $x_1x_2x_3x_4$  is equal to

- (a) 0
- (b) 1
- (c) 2
- (d) 3
- (e) Not enough information

Please turn over






I. Consider the nine lines  $y = 2x + 1, y = 2x + 2, \dots, y = 2x + 9$  and the seven lines  $y = -x + 1, y = -x + 2, \dots, y = -x + 7$ .

How many distinct points are there at which the line  $y = 1 - 10x$  crosses one or more of the other lines?

- (a) 12
- (b) 13
- (c) 14
- (d) 15
- (e) 16

Please turn over

**J.** Which of the following is the graph of  $y(y^3 + 4y^2x + 4x^3) = x^2(1 - x^2 - 6y^2)$ ?



End of last question