



CAMBRIDGE ASSESSMENT

STEP Solutions 2008

Mathematics

STEP 9465, 9470, 9475

STEP III, Solutions
June 2008

STEP Mathematics III 2008: Solutions

1. Following the hint yields

$$ax^2 + by^2 + (a+b)xy = \frac{1}{3}(x+y)$$

$$\text{which is } \frac{1}{5} + xy = \frac{1}{3}(x+y)$$

The same trick applied to the third equation gives $\frac{1}{7} + \frac{1}{3}xy = \frac{1}{5}(x+y)$.

The two equations can be solved simultaneously for xy and $(x+y)$, giving

$$xy = \frac{3}{35} \text{ and } (x+y) = \frac{6}{7}$$

Thus x and y are the roots of the quadratic equation $35z^2 - 30z + 3 = 0$ (x and y are interchangeable).

a and b are then found by substituting back into two of the original equations and the full solution is

$$x = \frac{3}{7} \pm \frac{2}{35}\sqrt{30} = \frac{3}{7} \pm \frac{2}{7}\sqrt{\frac{6}{5}}$$

$$y = \frac{3}{7} \mp \frac{2}{35}\sqrt{30} = \frac{3}{7} \mp \frac{2}{7}\sqrt{\frac{6}{5}}$$

$$a = \frac{1}{2} \mp \frac{\sqrt{30}}{36} = \frac{1}{2} \mp \frac{1}{6}\sqrt{\frac{5}{6}}$$

$$b = \frac{1}{2} \pm \frac{\sqrt{30}}{36} = \frac{1}{2} \pm \frac{1}{6}\sqrt{\frac{5}{6}}$$

2. (i) On the one hand

$$\sum_{r=0}^n [(r+1)^k - r^k] = \sum_{r=0}^n (r+1)^k - \sum_{r=0}^n r^k = \sum_{r=1}^{n+1} r^k - \sum_{r=0}^n r^k = (n+1)^k \text{ whilst expanding}$$

binomially yields

$$\begin{aligned} & k \sum_{r=0}^n r^{k-1} + \binom{k}{2} \sum_{r=0}^n r^{k-2} + \binom{k}{3} \sum_{r=0}^n r^{k-3} + \dots + \binom{k}{k-1} \sum_{r=0}^n r^{k-1} + \sum_{r=0}^n 1 \\ & = kS_{k-1}(n) + \binom{k}{2}S_{k-2}(n) + \binom{k}{3}S_{k-3}(n) + \dots + \binom{k}{k-1}S_1(n) + (n+1) \end{aligned}$$

and hence the required result.

Applying this in the case $k = 4$ gives

$$4S_3(n) = (n+1)^4 - (n+1) - \binom{4}{2}S_2(n) - \binom{4}{3}S_1(n)$$

which, after substitution of the two given results and factorization, yields the familiar

$$S_3(n) = \frac{1}{4}n^2(n+1)^2$$

The identical process with $k = 5$ results in

$$S_4(n) = \frac{1}{30}n(n+1)(6n^3 + 9n^2 + n - 1) = \frac{1}{30}n(n+1)(2n+1)(3n^2 + 3n - 1)$$

(ii) Applying induction, with the assumption that $S_t(n)$ is a polynomial of degree $t + 1$ in n for $t < r$ for some r , and then considering (*),

$(n+1)^{r+1} - (n+1)$ is a polynomial of degree $r + 1$ in n ,

and each of the terms $-\binom{r+1}{j}S_{r+1-j}(n)$ is a polynomial of degree $r + 2 - j$ in

n where $j \geq 2$, i.e. the degree is $\leq r$. A sum of polynomials of degree $\leq r + 1$ in n , is a polynomial of degree $\leq r + 1$ in n , and there is a single non-zero term in n^{r+1} from just $(n+1)^{r+1}$ so the degree of the polynomial is not reduced to $< r + 1$, i.e. it is $r + 1$. (The initial case is true to complete the proof.)

If, $S_k(n) = \sum_{i=0}^{k+1} a_i n^i = \sum_{r=0}^n r^k$ then $S_k(0) = a_0 + \sum_{i=1}^{k+1} a_i 0^i = \sum_{r=0}^0 r^k = 0$ and so $a_0 = 0$

$S_k(1) = \sum_{i=0}^{k+1} a_i 1^i = \sum_{r=0}^1 r^k = 1$ and so $\sum_{i=0}^{k+1} a_i = 1$ as required.

$$3. \quad \frac{dy}{dx} = \frac{b \cos \theta}{-a \sin \theta}$$

So the line ON is $y = \frac{a \sin \theta}{b \cos \theta} x$

SP is $y = \frac{b \sin \theta}{a(\cos \theta + e)}(x + ae)$

Solving simultaneously by substituting for x to find the y coordinate of T ,

$$y = \frac{b \sin \theta}{a(\cos \theta + e)} \left(\frac{b \cos \theta}{a \sin \theta} y + ae \right)$$

and using $b^2 = a^2(1 - e^2)$ to eliminate a^2 gives the required result.

Then the x coordinate of T is $\frac{b^2 \cos \theta}{a(1 + e \cos \theta)}$.


Eliminating θ using $\sec \theta + e = \frac{b^2}{ax}$ and $\tan \theta = \frac{by}{ax}$,

$$(x, y) \text{ satisfies } \left(\frac{b^2}{ax} - e \right)^2 = 1 + \left(\frac{by}{ax} \right)^2$$

and again using $b^2 = a^2(1 - e^2)$, this time to eliminate b^2 , gives, following simplifying algebra

$$(x + ae)^2 + y^2 = a^2, \text{ as required.}$$

4. (i)



The graph of $z = y$ has gradient 1 and passes through the origin.

The graph of $z = \tanh\left(\frac{y}{2}\right)$ which has gradient $\frac{1}{2} \operatorname{sech}^2\left(\frac{y}{2}\right) \leq \frac{1}{2}$ for $y \geq 0$ also passes through the origin and is asymptotic to $z = 1$.

Thus $y \geq \tanh\left(\frac{y}{2}\right)$ for $y \geq 0$.

$$\text{If } x = \cosh y, \text{ then } \sqrt{\frac{x-1}{x+1}} = \sqrt{\frac{\cosh y - 1}{\cosh y + 1}} = \sqrt{\frac{2 \sinh^2\left(\frac{y}{2}\right)}{2 \cosh^2\left(\frac{y}{2}\right)}} = \tanh\left(\frac{y}{2}\right)$$

and as $y \geq \tanh\left(\frac{y}{2}\right)$ for $y \geq 0$, $\operatorname{ar} \cosh x \geq \sqrt{\frac{x-1}{x+1}}$ for $x \geq 1$.

$$\sqrt{\frac{x-1}{x+1}} = \sqrt{\frac{x-1}{x+1}} \sqrt{\frac{x-1}{x-1}} = \frac{x-1}{\sqrt{x^2-1}} \text{ for } x > 1, \text{ and (*) is obtained.}$$

$$(ii) \quad \text{By parts } \int \operatorname{ar} \cosh x dx = x \operatorname{ar} \cosh x - \sqrt{x^2-1} + c$$

$$\text{and } \int \frac{x-1}{\sqrt{x^2-1}} dx = \sqrt{x^2-1} - \operatorname{ar} \cosh x + c'$$

$$\text{Thus } \int_1^x \operatorname{ar} \cosh x dx \geq \int_1^x \frac{x-1}{\sqrt{x^2-1}} dx \text{ for } x > 1 \text{ gives}$$

$$x \operatorname{ar} \cosh x - \sqrt{x^2-1} \geq \sqrt{x^2-1} - \operatorname{ar} \cosh x \text{ for } x > 1, \text{ which rearranges to give result}$$

(iii) Integrating (ii) similarly gives $x \operatorname{ar} \cosh x - \sqrt{x^2-1} \geq 2(\sqrt{x^2-1} - \operatorname{ar} \cosh x)$ for $x > 1$, which also can be rearranged as desired.

5. There are a number of correct routes to proving the induction, though the simplest is to consider $\left((T_{k+1}(x))^2 - T_k(x)T_{k+2}(x)\right) - \left((T_k(x))^2 - T_{k-1}(x)T_{k+1}(x)\right)$

For $f(x) = 0$, $(T_n(x))^2 - T_{n-1}(x)T_{n+1}(x) = 0$

and so $\frac{T_{n+1}(x)}{T_n(x)} = \frac{T_n(x)}{T_{n-1}(x)}$ provided that neither denominator is zero, leading to

$$\frac{T_n(x)}{T_{n-1}(x)} = \frac{T_1(x)}{T_0(x)} = r(x),$$

and so $\frac{T_n(x)}{T_{n-1}(x)} \times \frac{T_{n-1}(x)}{T_{n-2}(x)} \times \dots \times \frac{T_1(x)}{T_0(x)} = (r(x))^n$

Thus $T_n(x) = (r(x))^n T_0(x)$

Substituting this result into (*) for $n = 1$,

$\left((r(x))^2 - 2xr(x) + 1\right)T_0(x) = 0$, and as $T_0(x) \neq 0$, solving the quadratic gives

$$r(x) = x \pm \sqrt{x^2 - 1}$$

6. (i) Differentiating $y = p^2 + 2xp$ with respect to x gives

$$p = 2p \frac{dp}{dx} + 2x \frac{dp}{dx} + 2p \text{ which can be rearranged suitably.}$$

The differential equation $\frac{dx}{dp} + \frac{2}{p}x = -2$ has an integrating factor p^2 and integrating will give the required general solution.

Substituting $x = 2$, $p = -3$, leads to $A = 0$, i.e. $p = -\frac{3}{2}x$ which can be

substituted in the original equation and so $y = -\frac{3}{4}x^2$.

(ii) The same approach as in part (i) generates $\frac{dx}{dp} + \frac{2}{p}x = -\frac{(\ln p + 1)}{p}$,

which with the same integrating factor has general solution

$$x = -\frac{1}{4} - \frac{1}{2} \ln p + Bp^{-2}$$

and particular solution

$$x = -\frac{1}{2} \ln p - \frac{1}{4}$$


Again, substitution of $\ln p$ (and p) in the original equation leads to the solution

which is $y = -\frac{1}{2}e^{-2x-\frac{1}{2}}$

7. The starting point $c - a = \frac{1}{2}(1 + i\sqrt{3})(b - a)$ leads to the given result.

Interchanging a and b gives $2c = (a + b) + i\sqrt{3}(a - b)$ if A, B, C are described clockwise.

- (i) The clue to this is the phrase “can be chosen” and a sketch demonstrates that a pair of the equilateral triangles need to be clockwise, and the other pair anti-clockwise



Applying the results in the stem of the question to this configuration,

$$2p = (d + e) + i\sqrt{3}(e - d)$$

$$2q = (e + f) + i\sqrt{3}(e - f)$$

$$2r = (f + g) + i\sqrt{3}(g - f)$$

$$2s = (g + d) + i\sqrt{3}(g - d)$$

and so $2PS = (g - e) + i\sqrt{3}(g - e) = -2RQ$, PSQR is a parallelogram.

(The pairs could have been chosen with opposite parity leading to very similar working.)

- (ii) Supposing LMN is clockwise, U is the centroid of equilateral triangle LMH, V of MNJ, and W of NLK, then

$$3u = l + m + h \text{ where } 2h = (l + m) + i\sqrt{3}(m - l) \text{ with similar results for } v \text{ and } w.$$

Both $6w$, and $3[(u + v) + i\sqrt{3}(u - v)]$ can be shown to equal $3(n + l) + i\sqrt{3}(l - n)$ and so UVW is a clockwise equilateral triangle.

8. (i) $p = -\frac{1}{2}$

$$(1 + px)S = \frac{1}{3}x \text{ with all other terms cancelling and so } S = \frac{1}{3}x / \left(1 - \frac{1}{2}x\right) = \frac{2x}{3(2 - x)}$$

Using the sum of a GP

$$S_{n+1} = \frac{1}{3}x + \frac{1}{6}x^2 + \frac{1}{12}x^3 + \dots + \frac{1}{3 \times 2^n}x^{n+1} = \frac{\frac{1}{3}x \left(1 - \frac{x^{n+1}}{2^{n+1}}\right)}{\left(1 - \frac{x}{2}\right)}$$

Alternatively $S_{n+1} = S - (a_{n+2}x^{n+2} + \dots) = S - \frac{1}{2^{n+1}}x^{n+1}S$

$$= \left(1 - \frac{x^{n+1}}{2^{n+1}}\right) \frac{2x}{3(2-x)}$$

(ii) Using similar working to part (i)

$$18 + 8p + 2q = 0$$

$$37 + 18p + 8q = 0$$

$$\text{so } p = -\frac{5}{2}, q = 1$$

$$\text{and so } (1 + px + qx^2)T = 2 + 3x$$

$$\text{giving } T = (2 + 3x) \left/ \left(1 - \frac{5}{2}x + x^2\right) \right. = \frac{4 + 6x}{2 - 5x + 2x^2} = \frac{4 + 6x}{(2-x)(1-2x)}$$

$$\text{By partial fractions } T = \frac{14}{3}(1-2x)^{-1} - \frac{8}{3}\left(1 - \frac{x}{2}\right)^{-1}$$

$$\text{and so } T_{n+1} = \frac{14}{3}\left(1 + 2x + (2x)^2 + \dots + (2x)^n\right) - \frac{8}{3}\left(1 + \frac{x}{2} + \left(\frac{x}{2}\right)^2 + \dots + \left(\frac{x}{2}\right)^n\right)$$

$$= \frac{14}{3} \frac{1 - (2x)^{n+1}}{1 - 2x} - \frac{8}{3} \frac{\left(1 - \left(\frac{x}{2}\right)^{n+1}\right)}{1 - \frac{x}{2}}$$


Section B: Mechanics

9. When the particle starts to move, friction is limiting and so

$$mg \sin \pi T_0 - \mu mg = 0$$

$$\text{i.e. } \mu = \sin \pi T_0$$

(i)




When the particle comes to rest, the area under the acceleration-time graph is zero

$$\text{i.e. } \int_{T_0}^1 g \sin \pi t - \mu_0 g dt = 0$$

Completing the manipulation and eliminating T_0 using the relation at the start of the question renders the required result.

(ii)



In the case $\mu = \mu_0$, the motion is periodic with period 2, the particle is stationary in intervals $(0, T_0)$, $(1, 1 + T_0)$, $(2, 2 + T_0)$... , reversing its direction of motion after times 1, 2, 3, ... , and returning to its starting point at time 2 (and 4, 6, ...)

In the case $\mu = 0$, the motion is simple harmonic motion (period 2) superimposed on uniform motion, the particle instantaneously comes to rest at time 2, 4, ... but otherwise always moves in the positive x direction.

$$(x = \frac{g}{\pi^2}(\pi t - \sin \pi t))$$

10. Considering the r th short string $T_r = mg + T_{r-1}$

Also we have $T_r = \frac{\lambda x_r}{l}$, and $T_1 = mg$

$$\begin{aligned} \text{Thus } T_r &= rmg \text{ and so the total length is given by } \sum_1^n (l + x_r) = \sum_1^n \left(l + \frac{rmgl}{\lambda} \right) \\ &= nl + \frac{mgl}{\lambda} \frac{n(n+1)}{2} \end{aligned}$$

$$\text{The elastic energy stored is } \sum_1^n \frac{\lambda x_r^2}{2l} = \sum_1^n \lambda \left(\frac{lmg}{\lambda} \right)^2 \frac{r^2}{2l} = \frac{m^2 g^2 l}{12\lambda} n(n+1)(2n+1)$$

For the uniform heavy rope, we let $M = nm$, $L_0 = nl$, and consider the limit as $n \rightarrow \infty$

$$L = \lim \left(L_0 + \frac{M}{n} \frac{g}{\lambda} \frac{L_0}{n} \frac{n(n+1)}{2} \right) = L_0 \left(1 + \frac{Mg}{2\lambda} \right)$$

and the elastic energy stored is

$$\lim \left(\frac{m^2 g^2 l}{12\lambda} n(n+1)(2n+1) \right) = \lim \left(\frac{M^2 g^2 L_0}{12\lambda} \frac{n(n+1)(2n+1)}{n^3} \right) = \frac{M^2 g^2 L_0}{6\lambda}$$

and eliminating M using the result just found for L we obtain $\frac{2\lambda(L - L_0)^2}{3L_0}$

11. If the resistance couple (constant) is L , then using $L = I\alpha$ for the second phase of the motion, $L = \frac{I\omega_0}{T}$ and rotational kinetic energy used up doing work against the couple in the second phase gives

$$\frac{1}{2} I\omega_0^2 = L \times n_2 \times 2\pi$$

Hence, eliminating L and simplifying gives the first result.

If the particle descends a distance h in the first phase of motion, then $h = 2\pi r n_1$. If the particle has speed v at the end of the first phase, then $v = r\omega_0$


and using the work-energy principle,

$$mgh - L \times n_1 \times 2\pi = \frac{1}{2} I\omega_0^2 + \frac{1}{2} mv^2$$

Hence, eliminating h , v and ω_0^2 obtains the second result.

Section C: Probability and Statistics

12.



$$\begin{aligned}
 M_x(\theta) &= \int_{-\infty}^{\infty} e^{\theta x} f(x) dx = \int_{-\infty}^0 \frac{1}{2} e^{x(1+\theta)} dx + \int_0^{\infty} \frac{1}{2} e^{-x(1-\theta)} dx \\
 &= \frac{1}{2(1+\theta)} \left[e^{x(1+\theta)} \right]_{-\infty}^0 - \frac{1}{2(1-\theta)} \left[e^{-x(1-\theta)} \right]_0^{\infty} = \frac{1}{2(1+\theta)} + \frac{1}{2(1-\theta)} \quad (\text{requiring } |\theta| < 1) \\
 &= (1-\theta^2)^{-1}
 \end{aligned}$$

$$\begin{aligned}
 \text{Var}(X) &= M_x''(0) - (M_x'(0))^2 \\
 M_x'(\theta) &= 2\theta(1-\theta^2)^{-2}, \quad M_x''(\theta) = 2(1-\theta^2)^{-2} + 8\theta(1-\theta^2)^{-3} \\
 \text{and so } M_x'(0) &= 0, \quad M_x''(0) = 2, \quad \text{Var}(X) = 2
 \end{aligned}$$

$$\begin{aligned}
 \text{Or alternatively, } M_x(\theta) &= E(e^{\theta X}) = E\left(1 + \theta X + \frac{1}{2}\theta^2 X^2 + \dots\right) \\
 &= (1-\theta^2)^{-1} = 1 + \theta^2 + \theta^4 + \dots
 \end{aligned}$$

$$\text{and so } E(X) = 0, \quad E\left(\frac{1}{2} X^2\right) = 1, \quad \text{Var}(X) = E(X^2) - (E(X))^2 = 2$$

$$\text{If } T = Y/\sqrt{2n}, \text{ then } M_T(\theta) = E(e^{\theta T}) = E\left(e^{\theta \sum (x_i/\sqrt{2n})}\right) = \prod_{i=1}^n E\left(e^{\frac{\theta}{\sqrt{2n}} x_i}\right) = \left(1 - \frac{\theta^2}{2n}\right)^{-n}$$

$$\log(M_T(\theta)) = -n \log\left(1 - \frac{\theta^2}{2n}\right) = -n \left[-\frac{\theta^2}{2n} - \frac{\theta^4}{8n^2} - \frac{\theta^6}{24n^3} - \dots \right] = \frac{\theta^2}{2} + \frac{\theta^4}{8n} + \frac{\theta^6}{24n^2} + \dots$$

Thus as $n \rightarrow \infty$, $\log(M_T(\theta)) \rightarrow \frac{\theta^2}{2}$, and so $M_T(\theta) \rightarrow \exp\left(\frac{\theta^2}{2}\right)$

$$P(|Y| \geq 25) = 0.05 \text{ and } P\left(|Y/\sqrt{2n}| \geq 1.96\right) = 0.05 \text{ and so}$$

$$25 = 1.96\sqrt{2n}$$

$$2n = \frac{25^2}{1.96^2} \approx \frac{625}{4}$$

$$n \approx \frac{625}{8} \approx 78$$

$$13. \quad P(\text{1 ring created at first step}) = \frac{1}{2n-1},$$

$$P(\text{0 rings created at first step}) = \frac{2n-2}{2n-1}$$

$$E(\text{number of rings created at first step}) = \frac{1}{2n-1} \times 1 + \frac{2n-2}{2n-1} \times 0 = \frac{1}{2n-1}$$

Regardless of what happens at first step, after the first step there $2n-2$ free ends. Similarly after second step $2n-4$ free ends regardless, etc.

$$E(\text{number of rings at end of process}) = \frac{1}{2n-1} + \frac{1}{2n-3} + \frac{1}{2n-5} + \frac{1}{2n-7} + \dots + \frac{1}{1}$$

$\text{Var}(\text{number of rings at end of process}) =$

$$\frac{1}{2n-1} - \left(\frac{1}{2n-1}\right)^2 + \frac{1}{2n-3} - \left(\frac{1}{2n-3}\right)^2 + \frac{1}{2n-5} - \left(\frac{1}{2n-5}\right)^2 + \frac{1}{2n-7} - \left(\frac{1}{2n-7}\right)^2 + \dots + \frac{1}{1} - \left(\frac{1}{1}\right)^2$$

(as numbers of rings created at each step are independent)

$$= \frac{2(n-1)}{(2n-1)^2} + \frac{2(n-2)}{(2n-3)^2} + \frac{2(n-3)}{(2n-5)^2} + \dots + \frac{2}{3^2}$$

$$\text{For } n = 40000, E(\text{number of rings created}) = 1 + \frac{1}{3} + \frac{1}{5} + \dots + \frac{1}{79999}$$

$$= 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{80000} - \left(\frac{1}{2} + \frac{1}{4} + \dots + \frac{1}{80000}\right)$$

$$\approx \ln 80000 - \frac{1}{2} \ln 40000$$

$$= 2 \ln 20$$

$$\approx 6$$